

Background

Prolactin (PRL) is a neuroendocrine pituitary hormone. Prolactin is synthesized by the anterior pituitary, placenta, brain, uterus, dermal fibroblasts, decidua, B cell, T cells, NK cells, and breast cancer cells. Originally characterized as a lactogenic hormone, studies have demonstrated broader roles in breast cancer development, regulation of reproductive function, and immunoregulation. In the immune system, prolactin has been shown to be secreted by human PBMC and to act as a proliferative growth factor. Additionally, prolactin treatment of human PBMC has been shown to enhance IFN- γ production. Prolactin has several molecular forms. The predominant form is a monomer, the non-glycosylated form is 23 kDa and the glycosylated form is 25 kDa. Glycosylated prolactin is removed from the circulation faster and has been reported to have lower biological potency. Prolactin cDNA encodes a 227 amino acid residue protein with a putative 28 aa residue signal peptide. The prolactin receptor is a transmembrane type I glycoprotein that belongs to the cytokine hematopoietic receptor family. B cells, T cells, macrophages, NK cells, monocytes, CD34⁺ progenitor cells, neutrophils, mammary gland, liver, kidney, adrenals, ovaries, testis, prostate, seminal vesicles, and hypothalamus have all been shown to express the prolactin receptor. Three forms of the receptor, generated by differential splicing, have been identified. These isoforms differ in the length of their cytoplasmic domains. It is believed that the short cytoplasmic form is non-functional. Prolactin signal transduction involves the JAK/STAT families and Src kinase family.

References:

1. Cooke, N.E. *et al.* (1981) *J. Biol. Chem.* **256**:4007.
2. Ben-Johnson, N. *et al.* (1996) *Endoc. Rev.* **17**:639.
3. Cesario, T. *et al.* (1994) *Proc. Soc. Exp. Biol. Med.* **205**:89.
4. Price, A.E. *et al.* (1995) *Endoc.* **136**:4827.
5. Hoffmann, T. *et al.* (1993) *J. Endoc. Invest.* **16**:807.
6. Bellone, G. *et al.* (1995) *J. Cell Physiol.* **163**:221.
7. Cole, E. *et al.* (1991) *Endoc.* **129**:2639.
8. Lewis, U. *et al.* (1985) *Endoc.* **116**:359.

Description

Source	<i>E. coli</i> -derived Leu29 - Cys227, with an N-terminal Met Accession # Q5THQ0
N-terminal Sequence Analysis	Met
Predicted Molecular Mass	24 kDa

Specifications

Activity	Measured in a cell proliferation assay using Nb2-11 rat lymphoma cells. Gout, P.W. <i>et al.</i> (1980) <i>Cancer Res.</i> 40 :2433. The ED ₅₀ for this effect is typically 0.03 - 0.1 ng/mL.
Endotoxin Level	<1.0 EU per 1 μ g of the protein by the LAL method.
Purity	>97%, by SDS-PAGE under reducing conditions and visualized by silver stain.
Formulation	Lyophilized from a 0.2 μ m filtered solution in Acetonitrile and TFA. See Certificate of Analysis for details.

Preparation and Storage

Reconstitution	Reconstitute at 100 μ g/mL in sterile 4 mM HCl containing 1 mg/mL bovine serum albumin.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 3 months, -20 to -70 °C under sterile conditions after reconstitution.

12/11/2009

FOR RESEARCH USE ONLY
NOT FOR USE IN HUMANS