

DESCRIPTION



Species Reactivity	Human/Mouse/Rat
Specificity	Detects human, mouse, and rat FKBP51. In Western blots, less than 1% cross-reactivity with recombinant human FKBP12, FKBP13, FKBP38, and FKBP52 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human FKBP51 Thr2-Val457 Accession # Q13451
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.3 µg/mL	See Below
Immunohistochemistry	5-15 µg/mL	See Below

DATA

Western Blot	Immunohistochemistry
 <p>Detection of Human/Mouse/Rat FKBP51 by Western Blot. Western blot shows lysates of DU145 human prostate carcinoma cell line. PVDF membrane was probed with 0.3 µg/mL of Human/Mouse/Rat FKBP51 Antigen Affinity-purified Polyclonal Antibody (Catalog # AF4094) followed by HRP-conjugated Anti-Goat IgG Secondary Antibody (Catalog # HAF109). A specific band was detected for FKBP51 at approximately 51 kDa (as indicated). This experiment was conducted under reducing conditions and using Immunoblot Buffer Group 2.</p>	 <p>FKBP51 in Human Prostate. FKBP51 was detected in immersion fixed paraffin-embedded sections of human prostate using 15 µg/mL Human/Mouse/Rat FKBP51 Antigen Affinity-purified Polyclonal Antibody (Catalog # AF4094) overnight at 4 °C. Tissue was stained with the Anti-Goat HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # CTS008) and counterstained with hematoxylin (blue). View our protocol for Chromogenic IHC Staining of Paraffin-embedded Tissue Sections.</p>

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month from date of receipt, 2 to 8 °C, reconstituted. ● 6 months from date of receipt, -20 to -70 °C, reconstituted.

BACKGROUND

FK506 binding protein, 51 kDa molecular weight (FKBP51) is a peptidyl-prolyl isomerase that catalyzes the transition between *cis*- and *trans*- proline residues critical for proper folding of proteins. The macrolide immunosuppressants FK506 and Rapamycin are FKBP51 inhibitors. FKBP51 levels are induced by glucocorticoids. It associates with HSP90 complexes that are critical for the proper folding of steroid receptors. Single nucleotide polymorphisms in FKBP51 have been associated with major depression and hyper-responsiveness to antidepressants.