

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human P4HB in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human P4HB Asp18-Lys505 Accession # P07237
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 µg/mL	Recombinant Human Protein Disulfide Isomerase/P4HB (Catalog # 4236-DI)
Immunoprecipitation	25 µg/mL	Conditioned cell culture medium spiked with Recombinant Human Protein Disulfide Isomerase/P4HB (Catalog # 4236-DI), see our available Western blot detection antibodies

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month from date of receipt, 2 to 8 °C, reconstituted. ● 6 months from date of receipt, -20 to -70 °C, reconstituted.

BACKGROUND

P4HB (Prolyl 4-hydroxylase beta chain; also PDI) is a 60 kDa member of the protein disulfide isomerase family. As an intracellular homodimer, it forms a tetrameric complex with P4H alpha chains to form an active prolyl 4 hydroxylase. This catalyses the hydroxylation of proline in collagen. On the cell surface, it reduces disulfide bonds in HIV that allow the virus to fuse with CXCR4 and enter susceptible cells. Mature human P4HB is 491 amino acids (aa) in length. It contains two TRX domains (aa 25 134 and 368 475) plus an ER retention sequence (aa 505 508). There is one potential isoform that shows an 11 aa substitution for the first 162 amino acids. Over aa 18 505, human P4HB shares 94% aa identity with mouse P4HB.