



Biotinylated **Anti-human GASP-1/WFKKNRP Antibody**

ORDERING INFORMATION

Catalog Number: BAF2070

Lot Number: URW01

Size: 50 µg

Formulation: 0.2 µm filtered solution in PBS
with BSA

Storage: -20° C

Reconstitution: sterile 0.1% BSA in TBS

Specificity: human GASP-1

Immunogen: NS0-derived rhGASP-1

Ig Type: goat IgG

Applications: Western blot
ELISA detection

Preparation

Produced in goats immunized with purified, NS0-derived, recombinant human Growth and differentiation factor-Associated Serum Protein-1 (rhGASP-1; aa 35 - 576). Human GASP-1 specific IgG was purified by human GASP-1 affinity chromatography and then biotinylated.

Formulation

Lyophilized from a 0.2 µm filtered solution in phosphate-buffered saline (PBS) containing 50 µg of bovine serum albumin (BSA) per 1 µg of antibody.

Reconstitution

Reconstitute with sterile Tris-buffered saline pH 7.3 (20 mM Trizma base, 150 mM NaCl) containing 0.1% BSA. If 1 mL of buffer is used, the antibody concentration will be 50 µg/mL.

Storage

Lyophilized samples are stable for twelve months from date of receipt when stored at -20° C to -70° C. Upon reconstitution, the antibody can be stored at 2° - 8° C for 1 month without detectable loss of activity. Reconstituted antibody can also be aliquotted and stored frozen at -20° C to -70° C **in a manual defrost freezer** for six months without detectable loss of activity. **Avoid repeated freeze-thaw cycles.**

Specificity

This antibody has been selected for use as a detection antibody in human GASP-1 ELISAs and Western blots.

Applications

Western blot - This antibody can be used at 0.1 - 0.2 µg/mL with the appropriate secondary reagents to detect human GASP-1. The detection limit for rhGASP-1 is approximately approximately 0.5 ng/lane and 2 ng/lane under non-reducing and reducing conditions, respectively.

ELISA detection - This biotinylated antibody can be used as a detection reagent in a human GASP-1 sandwich immunoassay in combination with the human GASP-1 capture reagent (Cat. # MAB20702) and recombinant human GASP-1 (Cat. # 2070-GS) as the standard. The suggested concentration range for this detection reagent is 0.1 - 0.4 µg/mL and should be titrated to determine the optimal concentration. A general protocol is provided at www.RnDSystems.com/MAPELISA. In this format, less than 0.1% cross-reactivity with rhGASP-2 is observed.

Optimal dilutions should be determined by each laboratory for each application.