

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human XCR1 in Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human XCR1 Met1-Thr31, Ser89-Lys103, His168-Val190, Phe251-Tyr267 Accession # P46094
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 µg/mL	Recombinant Human XCR1
Flow Cytometry	2.5 µg/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile Tris-buffered saline, pH 7.3 (20 mM Trizma base, 150 mM NaCl) containing 0.1% bovine serum albumin.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

XCR1, also known as GPR5 and lymphotactin/SCM-1 (single cysteine motif 1) receptor, is a 38 kDa member of the G-protein coupled receptor 1 family. It binds XCL1/lymphotactin/SCM-1 α and XCL2/SCM-1 β . In addition, human herpesvirus 8 (HHV8) encodes two viral chemokines vCCL2/vMIP-II and vCCL3/vMIP-III that function as an antagonist and a highly selective agonist, respectively, for XCR1. XCR1 is expressed on neutrophils, CD8⁺ T cells, NK cells, CD4⁺ T cells and B cells. Human XCR1 is a 333 amino acid (aa), 7-transmembrane molecule. It contains a 32 aa N-terminus that lacks glycosylation sites and a 42 aa C-terminal cytoplasmic tail. Over the extracellular regions used for immunization, human XCR1 shares 62%, 54% and 64% aa identity with canine, mouse and porcine XCR1, respectively.