

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human CXCR3.
Source	Monoclonal Mouse IgG ₁ Clone # 49801
Purification	Protein A or G purified from ascites
Immunogen	NS0 mouse myeloma cell line transfected with human CXCR3 Met1-Leu368 Accession # P49682
Endotoxin Level	<0.10 EU per 1 µg of the antibody by the LAL method.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS and NaCl with Trehalose. See Certificate of Analysis for details.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	2.5 µg/10 ⁶ cells	Human whole blood lymphocytes
Immunohistochemistry	8-25 µg/mL	See Below
Neutralization	Measured by its ability to neutralize CXCL11/I-TAC-induced chemotaxis in the BaF3 mouse pro-B cell line transfected with human CXCR3. The Neutralization Dose (ND ₅₀) is typically 0.3-1.5 µg/mL in the presence of 7 ng/mL Recombinant Human CXCL11/I-TAC.	

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

CXCR3 is a G protein-coupled chemokine receptor that binds the α chemokines MIG (CXCL9), IP-10 (CXCL10), and I-TAC (CXCL11). CXCR3 is expressed on activated T cells, B cells, and NK cells.